











Bulletin,  
Agricultural  
Research  
Instt,  
Pusa, BENGAL,  
1923-1927,  
Nos:-151-168,



Bulletin No. 151

1923

Agricultural Research Institute, Pusa

---

**Method for the Accurate Determination of  
Carbonic Acid present as Carbonate in Soils**

BY

**PHANI BHUSAN SANYAL, M.Sc.,**  
*Assistant to the Imperial Agricultural Chemist.*



CALCUTTA  
SUPERINTENDENT GOVERNMENT PRINTING, INDIA  
1099



# **Agents for the Sale of Books Published by the Superintendent of Government Printing, India, Calcutta.**

## **IN EUROPE.**

Constable & Co., 10, Orange Street, Leicester Square,  
London, W.C.  
Kegan Paul, Trench, Trübner & Co., 68-74, Carter Lane,  
E.C., and 89, New Oxford Street, London, W.C.  
Bernard Quaritch, 11, Grafton Street, New Bond Street,  
London, W.  
F. & S. King & Sons, 2 & 4, Great Smith Street, West-  
minster, London, S.W.  
H. S. King & Co., 65, Osnihill, E.C., and 9, Pall Mall,  
London, W.  
Grindlay & Co., 54, Parliament Street, London, S.W.  
Lusac & Co., 46, Great Russell Street, London, W.C.  
W. Thacker & Co., 2, Creed Lane, London, E.C.

T. Fisher Unwin, Ltd., 1, Adelphi Terrace, London,  
W.C.  
Wheldon & Wesley, Ltd., 2, 3 & 4, Arthur Street, New  
Oxford Street, London, W.C. 2.  
East and West, Ltd., 8, Victoria Street, London,  
S. W. 1.  
B. H. Blackwell, 50 & 51, Broad Street, Oxford.  
Deighton Bell & Co., Ltd., Cambridge.  
Oliver and Boyd, Twickenham Court, Edinburgh.  
E. Ponsonby, Ltd., 110, Grafton Street, Dublin.  
Ernest Leroux, 28, Rue Bonaparte, Paris.  
Martinus Nijhoff, The Hague, Holland.  
Otto Harrassowitz, Leipzig.  
Friedländer and Sohn, Berlin.

## **IN INDIA AND CEYLON.**

Thacker, Spink & Co., Calcutta and Shill.  
Newman & Co., Ltd., Calcutta.  
B. Cambrey & Co., Calcutta.  
S. K. Lahiri & Co., Calcutta.  
B. Banerjee & Co., Calcutta.  
The Indian School Supply Depot, 399, Bow Bazar Street,  
Calcutta, and 326, Nawabpur, Dacca.  
Butterworth & Co. (India), Ltd., Calcutta.  
Rai M. C. Sarcar Bahadur & Sons, 90/21, Harrison  
Road, Calcutta.  
The Welton Library, 26, Chowringhee Road, Calcutta.  
Standard Literature Co., Ltd., Calcutta.  
Lal Chand & Sons, Calcutta.  
Association Press, Calcutta.  
The International Buddhist Book Depot, 4, Chaudul  
Chowk 1st Lane, Calcutta.  
Higginbotham & Co., Madras.  
V. Kalyaswami Iyer & Co., Madras.  
G. A. Natesan & Co., Madras.  
S. Murthy & Co., Madras.  
Thompson & Co., Madras.  
Temple & Co., Madras.  
P. R. Rama Iyer & Co., Madras.  
Vas & Co., Madras.  
E. M. Gopalakrishna Kone, Madras.  
Thacker & Co., Ltd., Bombay.  
D. B. Taraporevala, Sons & Co., Bombay.  
Mrs. Radhabai Atmaram Sagoo, Bombay.  
Sunder Poudurang, Bombay.  
Gopal Narayan & Co., Bombay.  
Bam Chandra Govind & Son, Kalbadevi, Bombay.  
Proprietor, New Kitabkhana, Poona.

The Standard Bookstall, Karachi, Rawalpindi and Murree  
Mangaldas Markandandas, Surat.  
Karsandas Narandas & Sons, Surat.  
A. H. Wheeler & Co., Allahabad, Calcutta and Bombay.  
N. B. Mathur, Supt., Nasir Kannon Hindi Press,  
Allahabad.  
Munshi Soela Ram, Managing Proprietor, Indian Army  
Book Depot, Jubb, Cawnpore.  
Rai Sahib M. Gulab Singh & Sons, Musli-Ain Press,  
Lahore and Allahabad.  
Rama Krishna & Sons, Lahore.  
Oxford Book and Stationery Co., Delhi.  
Supt., American Baptist Mission Press, Rangoon.  
Proprietor, Rangoon Times Press, Rangoon.  
The Modern Publishing House, Ltd., 70, Sparks Street,  
Rangoon.  
Manager, the "Hitavada," Nagpur.  
S. C. Talukdar, Proprietor, Students and Company,  
Cooch Behar.  
A. M. & J. Ferguson, Ceylon.  
Manager, Educational Book Depot, Nagpur and  
Jubbulpore.\*  
Manager of the Imperial Book Depot, 63, Chaudney  
Chauk Street, Delhi.\*  
Manager, "The Agra Medical Hall and Co-operative  
Association, Ltd." (Successors to A. John & Co.,  
Agra).  
Supt., Basel Mission Book and Tract Depository,  
Mangalore.\*  
P. Varadachary & Co., Madras.\*  
Ram Dayal Agarwala, 134, Katra, Allahabad.\*  
D. C. Anand & Sons, Peshawar.\*  
Manager, Newal Kishore Press, Lucknow.\*  
Maung Lo Gale, Proprietor, Law Book Depot, Mandalay,  
Times of Ceylon Co., Ltd.†

\* Agents for the sale of Legislative Department publications only.

† Agents for the sale of Archaeological publications only.

## A Method for the Accurate Determination of Carbonic Acid present as Carbonate in Soils

(Received for publication on 22nd March 1922.)

The amount of organic carbon in a soil is usually determined by deducting the amount of carbon dioxide ( $\text{CO}_2$ ) present as mineral carbonate from the total carbon dioxide ( $\text{CO}_2$ ) obtained by its complete combustion and that obtained by the decomposition of the residue of combustion with acid. The amount of organic carbon in Pusa soil is very small compared to the amount present as carbonate, and consequently in order to obtain reliable values for the former it is necessary to determine the latter with great accuracy.

The main difficulty in the determination of  $\text{CO}_2$  of mineral carbonate in soil, as pointed out by F. S. Marr<sup>1</sup> and J. C. Schollenberger<sup>2</sup> and verified by a series of determinations carried out at Pusa [Expts. I, II & V], lies in the fact that, on boiling a soil with acid, the organic substances present in it are decomposed and give out  $\text{CO}_2$ . Marr also observed that  $\text{CO}_2$  is liberated when some soils are boiled with water under atmospheric pressure. Our experience of the decomposition of Pusa soil with dilute boiling acid shows that, in spite of the soil containing only a small amount of organic matter, the results obtained with about one gram of soil are not sufficiently accurate. [Expt. I (1), (2) and (3).] The error, however, is considerably minimised by taking a larger quantity of soil for decomposition, but the results thus obtained, in triplicate determinations, still do not agree very well one with the other [Expt. I (4) and (5)].

Because of these difficulties, Marr adopted the method of decomposing the soil carbonates by boiling with dilute acid under reduced pressure at  $50^\circ\text{C}$ . and absorbing the liberated gas ( $\text{CO}_2$ ) in standard alkali. This involves double titration which is somewhat objectionable.

Gravimetric estimation of  $\text{CO}_2$  by decomposing soil by Marr's method has been found to be impracticable, as under above conditions soda-lime tubes lose moisture, and sulphuric acid tubes give out spray owing to the rapid passage of gas through them on evacuation. By properly pre-

<sup>1</sup> *Jour. Agr. Sci.*, III (1908—1910).

<sup>2</sup> *Jour. Ind. Eng. Chem.*, VIII (1916), No. 5, pp. 427, 428.

paring the sulphuric acid tubes (described later on) formation of spray in the tubes may be prevented if the gas is aspirated under slightly reduced pressure. It is doubtful, however, whether this will be the case, and whether the  $\text{CO}_2$  will be completely absorbed, when the gas is drawn through the tubes in a rapid stream under the reduced pressure required to boil the dilute acid at  $50^\circ\text{C}$ . Moreover, a very complicated arrangement, such as a bath for constant temperature ( $50^\circ\text{C}$ .), an efficient Geysk or Töppler pump, and a reaction flask and tubes sufficiently strong to bear the pressure, is necessary for such an operation, and these generally are not available in ordinary laboratories.

Consequently an attempt was made to devise some method for the gravimetric determination of  $\text{CO}_2$  under atmospheric pressure and *without the employment of heat*. This is only possible if the soil carbonates could be completely decomposed by dilute acid in the cold and if by any means the dissolved  $\text{CO}_2$  could be completely liberated from the solution.

In Pusa soils the carbonates are present mostly as calcium carbonate and to a slight extent as magnesium carbonate. So, finely ground calcium carbonate and dolomite were qualitatively tested with dilute hydrochloric acid in the cold. The former was found to be completely dissolved and the latter left a slight residue of sand which was not further dissolved even on boiling with stronger acid. It was, therefore, assumed that the carbonates present in the soil would be completely decomposed by dilute hydrochloric acid *in the cold*.

Some experiments were next carried out to see whether the dissolved  $\text{CO}_2$  could be completely washed out by the evolution of hydrogen in the reaction mixture with aluminium and hydrochloric acid.

For this purpose sodium carbonate was decomposed in the cold with excess of hydrochloric acid and then hydrogen was generated by dropping in a piece of aluminium foil. The hydrogen was finally driven off from the tubes by aspirating  $\text{CO}_2$ -free air. In all cases it was found that if the hydrogen be generated at a proper rate it was possible to estimate all the  $\text{CO}_2$  liberated during the reaction either in the cold or when heat was employed [Expt. III A (1), (2) and III B (1), (2)].

The next question which arose was whether the presence of clay would occlude or adsorb any  $\text{CO}_2$  in the cold. To decide this point a quantity of kaolin, which had been previously treated with strong HCl and washed free of acid and dried, was mixed with some sodium carbonate. The mixture was decomposed with hydrochloric acid and then hydrogen was generated as before and  $\text{CO}_2$ -free air finally aspirated through the tubes to replace the hydrogen. The results obtained show that no occlusion or adsorption of  $\text{CO}_2$  took place [Expt. III A (3)].

sample of soil was next taken for the purpose of experiment, and here difficulty was experienced owing to the caking of the soil at the bottom of the flask leading to incomplete decomposition. This trouble was overcome by the use of a specially designed apparatus which contained a mercury-sealed stirring arrangement. Arrangements were also provided to allow of a piece of aluminium foil being introduced when necessary, *i.e.*, after the decomposition of the carbonates. The apparatus was so arranged that the acid could be introduced without coming in contact with the aluminium foil. A full description of the apparatus and the method of employment is given later.

A further difficulty was met owing to the adherence of the fine clay to the surface of the aluminium foil, diminishing the rate of the generation of hydrogen and so preventing the dissolved  $\text{CO}_2$  being driven off. This was overcome by the introduction, after the decomposition of the soil, of a few drops of copper-sulphate solution, thus forming an aluminium-copper couple and accelerating the generation of hydrogen. If mercury is substituted for copper-sulphate, the aluminium amalgam in presence of the acid generates hydrogen too rapidly to be controlled. The method of using copper-sulphate possesses the advantage that the evolution of hydrogen can be reduced by dilution, or, on the other hand, the reaction can be rendered more active by introducing further amounts of copper-sulphate as desired.

An apparatus and method were finally evolved by which the carbonates of the soil could be completely decomposed at the ordinary temperature with excess of dilute hydrochloric acid and the mixture stirred all the while. The  $\text{CO}_2$  which remains dissolved in the menstrum is driven out by a stream of hydrogen generated by the action of aluminium and hydrochloric acid and a few drops of copper-sulphate solution. At the end,  $\text{CO}_2$ -free air is drawn through the flask and the tubes by aspiration, to sweep away all the  $\text{CO}_2$  and finally to drive away all the hydrogen in the tubes.

Very good results (*i.e.*, triplicate determinations agreeing very well) have been obtained by the above method [Expt. IV (1), (2) (3)].

The results obtained by boiling the mixture of soil and acid for ten minutes after it is completely decomposed in the cold and then generating hydrogen for a longer time were only very slightly higher than those obtained in the cold [Expt. V (1), (2), (3)], showing that the decomposition of the mineral carbonates under the latter condition was complete. The small variations in the triplicate determinations made by boiling the soil with acid are probably due to the decomposition of the organic substances present.

**Description of the apparatus.**

The apparatus (Fig. 1) consists of a reaction flask (M) to which the stirring arrangements for the introduction of acid are attached (also shown separately in Fig. 2). The delivery tube of the flask is connected to a double condenser (P) which is attached to a "T"-piece provided with a three-way stopper (Q) at the junction of three arms. The other end of the "T"-piece is attached to the bulbous tube (S) containing a little  $\text{H}_2\text{SO}_4$ , which in turn is attached to three  $\text{H}_2\text{SO}_4$  tubes (T) of which the third one is weighed and regarded as the guard tube. Two soda-lime tubes (T') are employed to absorb the  $\text{CO}_2$  and two  $\text{H}_2\text{SO}_4$  tubes (T'') to retain the moisture given off by the soda-lime tubes. Finally there is a second bulbous tube (S') containing little sulphuric acid to show the rate of passage of the gas through the tubes. An aspirator is employed to draw the gases through the tubes.

The mercury-sealed stirrer and the acid introducing arrangements are shown in Figs. 2 and 3.




Fig. 2.




Fig. 3.




Fig. 1. Sketch of the apparatus used for the determination of CO<sub>2</sub> present as mineral carbonates in soil.

These consist of a glass rod (A) having a movable glass blade (B) attached to the flattened end of the rod. A bent copper wire (C) is fastened to the rod a little above B to detach the aluminium foil (AL) when required. The rod (A) is fixed into the bung (D) at the top and is surrounded by the tube (E), which passes through the bungs (G) and (H) and is widened somewhat at (F) to receive acid from the stoppered vessel (L). The outer jacket of the mercury-seal (J) is fitted to the bung (H) and the inner tube (K) fixed to the bung (D) and dips into the mercury in the outer jacket (J) of the mercury-seal. The stem of the stoppered funnel (I) passes through the bung (D) and is bent inward over the top end of the tube (F).

A soda-lime-tube (N) is attached to the top of the funnel (I) (Fig. 1), whenever required, for drawing  $\text{CO}_2$ -free air into the apparatus.

#### Method of procedure.

Five to six grams of finely ground soil are weighed and washed into the reaction vessel M with two lots of 5 c.c. each of  $\text{CO}_2$ -free water each time, and the apparatus fitted together.

A mixture of 10 c.c. of strong HCl and 5 c.c. water ( $\text{CO}_2$ -free) is introduced little by little from the stoppered funnel (I) and the soil stirred continuously by turning the bung (D), so that the evolution of gas may be regular and at a moderate rate, the rate being indicated in the first bulbed tube (S). When the whole of the acid is run in and no further quantities of gas are evolved on stirring, it is assumed that the decomposition of the soil carbonates is completed. The stirrer is now raised so that the piece of aluminium foil is displaced and falls into the acid and hydrogen begins to be evolved. [It will be seen from the sketch (Fig. 1) that the stirrer can be raised so much as to displace the aluminium foil without allowing any gas to escape through the mercury seal.]

As stated previously, the evolution of hydrogen in mixture of soil and dilute acid is slow and this is accelerated by introducing a few drops of  $\text{CuSO}_4$  solution, thus forming an aluminium-copper couple. Hydrogen is allowed to evolve at a moderate rate at first for 15 minutes and then, when the rate is slowing down, a further quantity of 10 c.c. of strong HCl is added and hydrogen is evolved for another 10 minutes.

At this stage the outer jacket (J) (Fig. 3) is filled with mercury, so that most of the gas ( $\text{CO}_2$ , if any) is driven out of the inner space of the mercury-seal arrangement; the acid funnel is next washed with a little  $\text{CO}_2$ -free distilled water to get rid of HCl fumes (which may afterwards come into the soda-lime tube). The soda-lime tube N is now attached and

a slow current of  $\text{CO}_2$ -free air is drawn in through the whole apparatus, by means of the aspirator, to sweep away any  $\text{CO}_2$  remaining. When this is completed, the connection of the reaction flask with the tubes is broken by turning the three-way tap (Q) and  $\text{CO}_2$ -free air aspirated through the absorption tubes for half-an-hour so as to displace any hydrogen from the U-tubes. Half an hour's aspiration was found to be amply sufficient to displace all the hydrogen. The weight of  $\text{CO}_2$  evolved is found by weighing the absorption tubes and the final drying tubes.

The rate of evolution of  $\text{CO}_2$  and hydrogen, and the passage of air into the tubes, should not exceed 1 litre per half-an-hour.

Some difficulty was experienced owing to the formation of spray in the  $\text{H}_2\text{SO}_4$  tubes but this can be prevented by preparing pumicee saturated with sulphuric acid in a special way\* and placing glass wool in the upper portions of the U-tubes.

The results obtained by this apparatus and procedure have proved to be very satisfactory. For Pusa soil, containing only a small amount of organic substance, the method of determining  $\text{CO}_2$  by decomposing the soil carbonates with boiling dilute hydrochloric acid gives fairly good results if a large amount of soil (5–7 grams) is taken for decomposition [Expt. 1 (4) and (5)]. But in the case of soil rich in organic matter, boiling under atmospheric pressure involves the danger of decomposing the unstable organic substances, and the new method is recommended in such cases.

### Summary and conclusions.

1. Carbonates in the calcareous Pusa soil can be completely decomposed with hydrochloric acid, without boiling, if the soil be efficiently stirred.
2. The soil occludes no carbonic acid under the conditions in which the experiments are carried out.
3. The dissolved  $\text{CO}_2$  in the acid liquid can be completely driven out by the generation of hydrogen in it.
4. The rate of evolution of hydrogen in the menstruum can be regulated by the introduction of copper-sulphate solution.
5. The  $\text{CO}_2$  of mineral carbonate in soil can be gravimetrically determined with great accuracy by the use of the special apparatus according to the method recommended.

---

\* Ignited pumice is boiled with water for half-an-hour. The water is next replaced with strong sulphuric acid and kept overnight. The sulphuric acid is changed three times. Prepared in this way the pumice will not float in strong  $\text{H}_2\text{SO}_4$ . It is then thoroughly drained and introduced into the tubes through the stem of a funnel.



## EXPERIMENTS.

	Quantity of soil taken	% C (Mineral)
I. Boiled for 5 minutes . . . . .	(1) 1 gram	3.855
	(2) "	3.822
	(3) "	3.905
	(4) 5--7 grams	3.860
	(5) "	3.873
II. Boiled for 1 minute—		
(a) H generated, not stirred internally	(1) 1--2 grams	3.794
	(2) "	3.816
	(3) "	3.861
	(4) "	3.741
	(5) "	3.825
	(6) "	3.794
	(7) "	3.809
	(8) "	3.846
	(9) "	3.806
(b) H generated, stirred internally	(1) About 6 grams	3.731
	(2) "	3.855
	(3) "	3.736
III. Decomposition of--		
(A) $\text{Na}_2\text{CO}_3$ in the cold and generation of H in the cold.	(1) 2--2.5 grams	11.243
	$\text{Na}_2\text{CO}_3$	
	(2) 2--2.5 grams	11.154
	$\text{Na}_2\text{CO}_3$	
	(3) 2--2.5 grams	11.120
(B) $\text{Na}_2\text{CO}_3$ boiled and H generated in hot solution.	$\text{Na}_2\text{CO}_3 + 3$ grams Kaolin	
	(1) 0.559 grams	11.310
	$\text{Na}_2\text{CO}_3$	
	(2) 0.5403 grams	11.200
	$\text{Na}_2\text{CO}_3$	
IV. Decomposition of soil in the cold by acid and liberation of $\text{CO}_2$ from the cold solution with H & $\text{CuSO}_4$ (few drops).		
	(1) 5--7 grams of soil	3.861
	(2) "	3.859
	(3) "	3.863
V. Decomposition of soil in the cold mixture, boiled afterwards for ten minutes and H generated for a long time.		
	(1) 5--7 grams of soil	3.889
	(2) "	3.865
	(3) "	3.876

## Publications of the Imperial Department of Agriculture in India.

To be had from :—

The Office of the Agricultural Adviser to the Government of India, Pusa, Bihar,  
and from the following Agents :—

- |  |   |
|--|---|
| (1) Thacker, Spink & Co., Calcutta.            | (7) Thacker & Co., Ltd., Bombay.              |
| (2) W. Newman & Co., Calcutta.                 | (8) Sunder Pandurang, Bombay.                 |
| (3) Rai M. C. Sarkar Bahadur & Sons, Calcutta. | (9) Rai Sahib M. Golab Singh & Sons, Lahore.  |
| (4) Higginbotham & Co., Madras.                | (10) Manager, Educational Book Depot, Nagpur. |
| (5) Thompson, Ltd., Madras.                    |   |
| (6) D. B. Tamporevala, Sons & Co., Bombay.     |   |

**A complete list of the publications of the Imperial Department of Agriculture in India can be obtained on application from the Agricultural Adviser to the Government of India, Pusa, Bihar, or from any of the above-mentioned agents.**

These publications are :—

1. *The Agricultural Journal of India.* A Journal dealing with subjects connected with agricultural economies, field and garden crops, economic plants and fruits, soils, manures, methods of cultivation, irrigation, climatic conditions, insect pests, fungus diseases, co-operative credit, agricultural cattle, farm implements, and other agricultural matters in India. Illustrations, including coloured plates, form a prominent feature of the Journal. It is edited by the Agricultural Adviser to the Government of India, and is issued once every two months or six times a year. *Annual Subscription*, Rs. 6 or 9s. 6d., including postage. *Single copy*, R. 1-8 or 2s.
2. Scientific Reports of the Agricultural Research Institute, Pusa.
3. Annual Review of Agricultural Operations in India.
4. Proceedings of the Board of Agriculture in India.
5. Proceedings of Sectional Meetings of the Board of Agriculture.
6. Memoirs of the Imperial Department of Agriculture in India :
  - (a) Botanical Series.
  - (b) Chemical Series.
  - (c) Entomological Series.
  - (d) Bacteriological Series.
  - (e) Veterinary Series.
7. Bulletins issued by the Agricultural Research Institute, Pusa.
8. Indigo Publications.
9. Books.

**The following are the publications of the last two years :—**

- Scientific Reports of the Agricultural Research Institute, Pusa (including the Reports of the Imperial Dairy Expert and the Secretary, Sugar Bureau), for the year 1920-21. Price, R. 1-8.
- Scientific Reports of the Agricultural Research Institute, Pusa (including the Reports of the Imperial Dairy Expert and the Secretary, Sugar Bureau), for the year 1921-22. Price As. 14.
- Review of Agricultural Operations in India, 1920-21. Price, R. 1-4.
- Review of Agricultural Operations in India, 1921-22. Price, R. 1-4.
- Proceedings of the Board of Agriculture in India, held at Pusa on the 13th February, 1922, and following days (with Appendices). Price, R. 1.
- Report of the Proceedings of the Fifth Entomological Meeting, held at Pusa on 5th—10th February, 1922. (*In the press*.)

## MEMOIRS OF THE DEPARTMENT OF AGRICULTURE IN INDIA.

### Botanical Series

- Vol. XI, No. VII. Correlation of Colour Characters in Rice, by G. P. HECTOR, M.A., D.Sc. Price, R. 1-4 or 1s. 8d.
- Vol. XI, No. VIII. The Inheritance of Characters in Rice, II, by F. R. PARNELL, M.A., AG. DIP. (Cantab.), with the assistance of G. N. RANGASWAMI AYYANGAR, B.A., K. RAMIAH, L.A.G., and C. R. SRINIVASA AYYANGAR, L.A.G. Price, R. 1-4 or 1s. 8d.
- Vol. XI, No. IX. A New Ginger Disease in Godavari District, by S. SUNDARAMAN, M.A. Price, R. 1 or 1s. 4d.
- Vol. XI, No. X. *Helminthosporium* spp. on Cereals and Sugarcane in India, Part I. Diseases of *Zea Mays* and *Sorghum vulgare* caused by species of *Helminthosporium*, by MASORANJAN MITRA, M.Sc. Price, R. 1 or 1s. 4d.
- Vol. XII, No. I. The Wheats of Bihar and Orissa, by ALBERT HOWARD, C.I.E., M.A., GABRIELLE L. C. HOWARD, M.A., and ABDUR RAHMAN KHAN. Price, As. 8 or 9d.
- Vol. XII, No. II. I—History of the operations against Bud-rot of Palms in South India. II—Inoculation Experiments with *Phytophthora palmivora* BURL. OR *Botryotinia doellingeri* Linn. and *Cocos necifera* Linn., by W. McRAE, M.A., B.Sc. Price, R. 1-4 or 2s.
- Vol. XII, No. III. Studies in Inheritance in Cotton. I. History of a Cross between *G. herbaceum* and *G. pygmaeum*, by G. L. KOTTER, M.A.G. Price, R. 1-4 or 1s. 9d.

### Chemical Series

- Vol. VI, No. I. Investigations on Indian Opium, No. 1. Non-environmental Factors influencing the Alkaloidal Content and Yield of Latex from the Opium Poppy (*Papaver somniferum*), by HAROLD E. ANNETT, D.Sc. (Lond.), F.I.C., M.S.E.A.C., HARI DAS SEN, M.Sc., and HAR DATAL SINGH, D.Sc. Price, R. 1-8 or 2s.
- Vol. VI, No. II. Investigations on Indian Opium, No. 2. The Effect of Non-environmental Factors on the Alkaloidal Content and Yield of Latex from the Opium Poppy (*Papaver somniferum*) and the bearings of the work on the functions of Alkaloids in Plant Life, by HAROLD E. ANNETT, D.Sc. (Lond.), F.I.C., M.S.E.A.C. Price, Rs. 2 or 2s. 6d.
- Vol. VI, No. III. Studies in Soil Moisture, Part I, by B. H. WILSDON, B.A. (Oxon.), I.E.S. Price, R. 1-8 or 2s.
- Vol. VI, No. IV & V. Variations in some Characteristics of the Fat of Buffalo and Cow Milk with changes in Season and Feeding; The Mutual Applicability of the Analytical Figure 8 for Butter Fat and Ghee, by F. J. FLYMEN, A.C.C.I., and A. R. PADMANABHA AYYAR, B.A. Price, As. 12 or 1s. 3d.
- Vol. VI, No. VI. Investigations on Indian Opium, No. 3. Studies in the Meconic Acid Content of Indian Opium, by HAROLD E. ANNETT, D.Sc. (Lond.), F.I.C., M.S.E.A.C., and MATURIA NATH BOSE, M.A. Price, As. 6 or 6d.
- Vol. VI, No. VII. Chemical Studies on Safflower Seed and its Germination, by V. A. TANHANE, M.Sc., M.A.L. Price, As. 10 or 1s.
- Vol. VI, No. VIII. Note on the Permanent Manurial Plots, Coimbatore, by ROLAND V. NORRIS, D.Sc. Price, R. 1-4 or 1s. 9d.
- Vol. VII, No. I. A Note on the Hydrocyanic Acid in the Burma Bean (*Phaseolus lunatus*, sp.), by F. J. WARREN, M.Sc., D.Sc. Price, As. 12 or 1s.
- Vol. VII, No. II. Studies of an Acid Soil in Assam, II, by A. A. MCGILL, D.Sc. Price, As. 12 or 1s.
- Vol. VII, No. III. A Preliminary Note on the Decomposition of Calcium Cyanamide in South Indian Soils, by ROLAND V. NORRIS, D.Sc., F.I.C., E. VISWANATH and C. V. RAMASWAMI AYYAR, L.A.G. (In the press.)

### Entomological Series

- Vol. VII, No. VI. Life-histories of Indian Insects. Diptera: *Sphyracephala learsiana*, Westw., by S. K. SEN, B.Sc. Price, As. 12 or 1s.
- Vol. VII, Nos. VII & VIII. New and Rare Indian Odonata in the Pusa Collection; Further Notes on *Rhinocypha* Larvae, by Major F. C. FRASER, I.M.S. Price, R. 1-4 or 1s. 9d.
- Vol. VII, No. IX. Notes on Indian Diptera, by RONALD SENIOR-WHITE, F.E.S. Price, R. 1-12 or 2s. 3d.
- Vol. VII, No. X. *Platyedra gossypiella*, Samml., Pink Boll-Worm in South India, 1920-21, by E. BALLARD, B.A., F.E.S. Price, As. 12 or 1s.
- Vol. VII, No. XI. Studies in Indian Dermaptera, by MORRIS HERARD. Price, R. 1-1 or 1s. 9d.
- Vol. VII, No. XII. Further Notes on *Pemphigus affinis*, Fsb. (The Cotton Stem Weevil), by E. BALLARD, B.A., F.E.S. Price, R. 1 or 1s. 4d.
- Vol. VII, No. XIII. An Account of Experiments on the Control of *Nipta* (*Schenophilus*) *incertellus* in the Golavadi Delta, by E. BALLARD, B.A., F.E.S. Price, As. 11 or 1s. 4d.
- Vol. VIII, Nos. I, II & III. Hydrophilidae of India, by A. DUCHEMINOT; An Annotated List of Ichneumonidae in the Pusa Collection, by G. R. DUFF, B.A.; A Second Note on Odonata in the Pusa Collection, by Major F. C. FRASER. Price, R. 1 or 1s. 6d.
- Vol. VIII, No. IV. Notes on Indian Muscids, by R. SENIOR-WHITE, F.E.S. Price As. 12 or 1s.

### Veterinary Series

- Vol. III, No. IV. Studies in Rinderpest, by W. A. POOL, M.R.C.V.S., and T. M. DOYLE, F.R.C.V.S. Price, R. 1 or 1s. 4d.
- Vol. III, No. V. Etiology of Equine Contagious Abortion in India, by T. M. DOYLE, F.R.C.V.S. Price, As. 8 or 9d.

### BULLETINS ISSUED BY THE AGRICULTURAL RESEARCH INSTITUTE,

#### PUSA

- No. 116. A Survey of the Indian Poppy-growing Districts for Morphine Content of the Opium produced, by H. E. ANNETT, D.Sc. (Lond.), F.R.C., M.S.E.A.C., HARI DAS SEN, M.Sc., and HAR DAYAL SINGH, B.Sc. Price, As. 2.
- No. 117. Experiments with Castor Seed Conducted at Sabour, by C. SOMERS TAYLOR, B.A. Price, As. 3.
- No. 118. The Saving of Irrigation Water in Wheat Growing, by ALBERT HOWARD, C.I.E., M.A., and GABRIELLE L. C. HOWARD, M.A. Price, As. 4.
- No. 119. The Agricultural Development of Baluchistan, by ALBERT HOWARD, C.I.E., M.A., and GABRIELLE L. C. HOWARD, M.A. Price, As. 6.
- No. 120. The Serum Simultaneous Method of Inoculation against Rinderpest, by W. A. POOL, M.R.C.V.S. Price As. 2. (*Order printed*)
- No. 121. Notes on Contagious Abortion in Pony and Donkey Mares, by R. BRANFORD, M.R.C.V.S., and T. M. DOYLE, F.R.C.V.S. Price, As. 5.
- No. 122. Pusa 12 and Pusa 4 in the Central Circle of the United Provinces, by B. C. DUTT, M.B.E., B.Sc., F.C.S., ALBERT HOWARD, C.I.E., M.A., and GABRIELLE L. C. HOWARD, M.A. Price, As. 11.
- No. 123. The Bundelkhand Cottons. Experiments in their Improvement by Pure Line Selection, by B. C. DUTT, M.B.E., B.Sc., F.C.S., and NIZAMUDDIN HANDEK. Price, As. 4.
- No. 124. Safflower Oil, by ALBERT HOWARD, C.I.E., M.A., and J. STEWART REMINGTON. Price, As. 4.

**BULLETINS ISSUED BY THE AGRICULTURAL RESEARCH INSTITUTE,  
PUSA—concd.**

- No. 125. The Weevil Fauna of South India with special reference to species of Economic Importance, by T. V. RAMAKRISHNA AYYAR, B.A., F.E.S., F.Z.S. Price, R. 1-4.
- No. 126. Cawnpore-American Cotton, II. Further Field Trials (1918-20), Spinning Trials and Market Organization, by B. C. BURT, M.B.E., B.Sc., F.R.S. Price, As. 4.
- No. 127. The Coconut Bleeding Disease, by S. SUNDARAMAN, M.A. Price, As. 8.
- No. 128. *Helminthosporium* Disease of Paddy, by S. SUNDARAMAN, M.A. Price, As. 9.
- No. 129. The Preparation of Anti-rinderpest Serum using Animals of moderate susceptibility as Virus Producers, by W. A. POOL, M.B.C.V.S., and T. M. DOYLE, F.R.C.V.S. Price, As. 12.
- No. 130. Feeding Experiments at Government Cattle Farm, Hissar, by R. BRANFORD, M.B.C.V.S., and E. SEWELL, M.C., M.B.C.V.S. Price, As. 3.
- No. 131. An Entomologist's Crop Pest Calendar for the Madras Presidency, by T. V. RAMAKRISHNA AYYAR, B.A., F.E.S., F.Z.S. Price, As. 2.
- No. 132. Report of Campaign against *Spodoptera mauritia*, Boisd. (*Noctuidæ*) in Malabar, by E. BALLARD, B.A., F.E.S. Price, As. 2.
- No. 133. Results of Investigation of Bionomics of *Platyedra gossypiella*, Saunders, in South India, together with some notes on *Earias insulana* and *E. fabia*, by E. BALLARD, B.A., F.E.S. Price, As. 5.
- No. 134. Supplementary Observations on Borers in Sugarcane, Rice, etc., by C. C. GHOSH, B.A., Price, As. 6.
- No. 135. Some Observations on the Control of Field Rats in the Punjab, by M. A. HUSSAIN, M.A., and HEM SINGH PRUTHI, M.Sc. Price, A. 1. (*Out of print*.)
- No. 136. The Hydrogen Ion Concentrations of some Indian Soils and Plant Juices, by W. R. G. ATKINS, O.B.E., Sc.D., F.R.S. Price, As. 4.
- No. 137. Note on the Probability of an Inter-relation between the Length of the Stigma and that of the Fibre in some forms of the genus *Gossypium*, by RAM PRASAD. Price, As. 4.
- No. 138. Methods of Examination of Certain Characters in Cotton, by G. R. HILSON, B.Sc. Price, As. 8.
- No. 139. List of Publications of Indian Entomology, 1920-21 (compiled by the Imperial Entomologist). Price, R. 1.
- No. 140. The Determination of Prussic Acid in Burma Beans (*Phaseolus lunatus*) (Preliminary Note), by J. CHARLTON, B.Sc. Price, As. 3.
- No. 141. Comparative Manurial Value of the whole Plants and the different parts of Green Manures, by N. V. JOSHI, B.A., M.Sc., L.Ag. Price, As. 6.
- No. 142. The Cultivation of Lac in the Plains of India, by C. S. MISRA, B.A. Price, R. 1-8.
- No. 143. Ber (*Zizyphus jujaba*) Fruit and its Fly Pest, by J. L. KHARE, F.E.S. Price, As. 6.
- No. 144. Observations on the Morphology and Life-cycle of *Filaria vesicula* Grassi, by M. ANANT NARAYAN RAO. Price, As. 6.
- No. 145. Some Observations on the Barren Soils of Lower Bari Doab Colony in the Punjab, by S. M. NASTR. Price, As. 3.
- No. 146. The Prevention of Nuisances caused by the Parboiling of Paddy, by J. CHARLTON, B.Sc., A.L.C. (*In the press*.)
- No. 147. List of Publications on Indian Entomology, 1922, compiled by the Imperial Entomologist. (*In the press*.)
- No. 148. The Relative Responsibility of Physical Heat and Micro-organisms for the Hot Weather Rotting of Potatoes in Western India, by S. L. ASHRAFI, B.A. (*In the press*.)
- No. 149. A Study of the Factors Operative in the value of Green Manure, by R. H. WILSON, B.A., LL.B., P. E. LANDER, M.A., B.Sc., A.L.C., and M. MURANDI LAL. (*In the press*.)
- No. 150. The Improvement of Fodder and Forage in India (Papers read before a joint meeting of the sections of Agriculture and Botany, Indian Science Congress, 1923), edited by GABRIELLE L. C. HOWARD, M.A. Price, As. 6.
- No. 151. A Method for the Accurate Determination of Carbonic Acid present as Carbonate in Soils, by PHANI BHUSAN SANYAL, M.Sc. Price, As. 2.

Journal No. 153

**Agricultural Research Institute, Pusa**

---

**Tamarind as a Source of Alcohol and Tartaric  
Acid**

BY

**H. N. BATHAM, M.A.,**  
*Offg. Agricultural Chemist, United Provinces*

AND

**L. S. NIGAM, L.Ag.,**  
*Assistant to Agricultural Chemist, United Provinces*



CALCUTTA : GOVERNMENT OF INDIA  
CENTRAL PUBLICATION BRANCH  
1924

Price As. 3 or 4d.



Agricultural Research Institute, Pusa

**Tamarind as a Source of Alcohol and Tartaric  
Acid**

BY

**H. N. BATHAM, M.A.,**

*Offg. Agricultural Chemist, United Provinces*

AND

**L. S. NIGAM, L.Ag.,**

*Assistant to Agricultural Chemist, United Provinces*



CALCUTTA : GOVERNMENT OF INDIA  
CENTRAL PUBLICATION BRANCH

1924



**Government of India publications are obtainable from the  
Government of India Central Publication Branch, 8, Hastings  
Street, Calcutta, and from the following agents.**

**IN EUROPE.**

Constable & Co., 10, Orange Street, Leicester Square,  
London, W.C.

Kegan Paul, Trench, Trübner & Co., 68-74, Carter  
Lane, E.C., and 39, New Oxford Street,  
London, W.C.

Bernard Quaritch, 11, Grafton Street, New Bond  
Street, London, W.

P. S. King & Sons, 2 & 4, Great Smith Street, West-  
minster, London, S.W.

Grindley & Co., 54, Parliament Street, London, S.W.

Luzac & Co., 46, Great Russell Street, London, W.C.

W. Thacker & Co., 2, Creed Lane, London, E.C.

T. Fisher Unwin, Ltd., 1, Adelphi Terrace, London,  
W.C.

Walden & Wesley, Ltd., 2, 3 and 4, Arthur Street,  
New Oxford Street, London, W.C. 2.

East and West, Ltd., 3, Victoria Street, London,  
S.W. 1.

B. H. Blackwell, 50 & 51, Broad Street, Oxford.

Deighton Bell & Co., Ltd., Cambridge.

Oli'er and Boyd, Tweeddale Court, Edinburgh.

K. Ponsonby, Ltd., 116, Grafton Street, Dublin.

Krnest Leroux, 28, Rue Bonaparte, Paris.

Martinus Nijhoff, The Hague, Holland.

Otto Harrassowitz, Leipzig.

Wiedländer und Sohn, Berlin.

**IN INDIA AND CEYLON.**

Thacker, Spink & Co., Calcutta and Simla.

W. Newman & Co., Ltd., Calcutta.

R. Cambay & Co., Calcutta.

S. K. Lahiri & Co., Calcutta.

The Indian School Supply Depot, 309, Bow Bazar  
Street, Calcutta, and 220, Nawabpur, Dacca.

Butterworth & Co. (India), Ltd., Calcutta.

Rai M. C. Sanyal Balador & Sons, 99/2-A, Harrison  
Road, Calcutta.

The Weldon Library, 26, Chowringhee Road,  
Calcutta.

Standard Literature Company, Limited, Calcutta.

Association Press, Calcutta.

Chakraverty, Chatterjee & Co., Ltd., 13, College  
Square, Calcutta.

Hazlebotham & Co., Madras.

V. Kalyanaram Iyer & Co., Madras.

P. R. Rama Iyer & Co., Madras.

Rehouse and Sons, Madras.

Bright & Co., Trivandrum.

V. S. Swaminathan, Bookseller, West Tower  
Street, Madras.

Thacker & Co., Ltd., Bombay.

D. B. Taraporevala, Sons & Co., Bombay.

Sunder Pandurang, Bombay.

Ram Chandra Govind & Sons, Kalladevi, Bombay.

Proprietor, New Kitabkhana, Poona.

The Manager, Oriental Book Supplying Agency,  
15, Shukrawar, Poona City.

The Standard Bookstall, Karachi, Rawalpindi,  
Murree, Lahore and Quetta.

Karnadas Narandas & Sons, Surat.

A. H. Wheeler & Co., Allahabad, Calcutta and  
Bombay.

N. B. Mathur, Supdt., Nazir Kanun Hind Press,  
Allahabad.

The North India Christian Tract and Book Society,  
18, Olive Road, Allahabad.

Ram Dayal Anurwala, 184, Katra, Allahabad.

Manager, Newal Kishore Press, Lucknow.

The Upper India Publishing House, Ltd., 41  
Aminabad Park, Lucknow.

Munshi Sactaram, Managing Proprietor, Indian  
Army Book Depot, Jubb, Cawnpore.

Rai Sahib M. Gulab Singh & Sons, Mudd-I-Am Press,  
Lahore and Allahabad.

Bans Krishna & Sons, Booksellers, Anarkali,  
Lahore.

Manager of the Imperial Book Depot, 68, Chandany  
Chauk Street, Delhi.

Oxford Book and Stationery Company, Delhi.

Supdt., American Baptist Mission Press, Rangoon.

Proprietor, Rangoon Times Press, Rangoon.

The Modern Publishing House, Ltd., 70, Sparks  
Street, Rangoon.

The International Indologist Book Depot, Post  
Box No. 971, Rangoon.

Manager, the "Hivavada," Nagpur.

S. C. Talukdar, Proprietor, Students & Co., Coonoor  
Bazaar.

Times of Ceylon Co., Ltd.

The Manager, Ceylon Observer, Colombo.

## Tamarind as a Source of Alcohol and Tartaric Acid.

(Received for publication on 7th January, 1924.)

India is full of resources, vegetable and mineral, and there are many raw products which are capable of being used for industrial purposes. The tamarind (*Tamarindus indica*), which is so widely found in India, has yet been only partially investigated, although every part of the tree is utilized for some purpose or other, the most important being the fruit. According to the varieties of fruit, the tree is divided into three classes :—(1) The sour fruited, (2) the sweet fruited and (3) the red fruited. The pulp of the fruit of the last variety is of a rose colour and commands the highest price and is preferred for preserving.

Of the three parts of the fruit, *viz.*, shell, seed and pulp, the last is the most useful. It is for the sake of the pulp that the fruits form a regular article of commerce. Besides their consumption for edible purposes in India, they are largely exported to Europe. According to "Tropical Agriculture,"<sup>1</sup> a new market has sprung up for tamarind in Italy, where it is in demand for the manufacture of syrups and sweetmeats. According to the same authority, the average annual yield of an adult tree is 350 lb. of pods or fruits, which represent 200 lb. of prepared pulp. The pulp is thus about 75 per cent. in the fruit. The average percentage of the seed and the pulp, as determined in the de-corticated fruits available in the Cawnpore market, is 32 and 68 respectively. Therefore, the greatest part of the fruit is the pulp, which principally consists of sugars and tartaric acid. The sugar content ranges from about 30 to 40 per cent. and tartaric acid from about 9 to 15 per cent. as can be seen from the analyses of different samples of tamarind procured from the Cawnpore market as well as from the analyses by other workers. The tartaric acid was determined by the revised form of Goldenburg method (1907),<sup>2</sup> and the reducing sugars by Söldani method modified by Ost, which was described in a Pusa Bulletin.<sup>3</sup> The results are given in Table I.

---

<sup>1</sup> *Tropical Agriculture*, Nov. 1905.

<sup>2</sup> Lunge and Keane. *Technical Methods of Analysis*.

<sup>3</sup> *Bull. 13 of Agri. Res. Inst. Pusa*.

TABLE I.

Showing the percentage of reducing sugars and tartaric acid in different samples of tamarind.

Sample No.	Reducing sugars	Tartaric acid
1	39.56	9.65
2	41.00	9.52
3	33.96	11.16
4	39.21	12.76
5	38.88	10.45
6	29.16	8.60
7	Traces	2.25
8	Appreciable quantity	5.51

There appears to be some relation between the contents of sugar and tartaric acid: the higher the sugar content the less the quantity of tartaric acid and *vice versa*. Sample No. 6 was one year old and seems to have deteriorated in its two contents as is the case with *mahua* (*Bassia latifolia*) flower when kept for a year or so, as mentioned later on. Samples Nos. 7 and 8 were unripe tamarinds in different stages, and it appears that probably both sugar and tartaric acid were in their formative stage.

It is due to the presence of these two constituents that the pulp is consumed by Indians in the form of congies, syrups, *chatni*, beverages, etc., but no industrial use has yet been made of it.

Some time ago, efforts were made in New Caledonia to distil alcohol and wine from banana and tamarind.<sup>1</sup> The latter produced an excellent brandy manufactured in the following way:—The fruit picked a little before it was ripe was crushed in water and the whole allowed to ferment. From 13.5 kilos of fruit 8 litres of brandy of good quality were produced. A very remarkable fact is the absence in the tamarind of the essential oils, which pollute almost all fruit or grain alcohols below a certain degree. Besides alcohol, there can be extracted from the fruit of the tamarind a certain quantity of tartaric acid.

Recently a process has been worked out by Dr. Sudborough and Mr. P. N. Védhachalam<sup>2</sup> to manufacture tartaric acid from the tamarind pulp. In this manufacture three difficulties had to be

<sup>1</sup> *Jour. Soc. Chem. Ind.*, May 1889.

<sup>2</sup> *Jour. Ind. Aust. Sci.*, III, 5.

encountered. The first, viz., that about the filtration of the aqueous extract of the pulp, was removed by the use of an autoclave. The second difficulty about the refining and bleaching the solution was solved by the use of different bleaching agents. The disposal of the bye-products was the third important point left for others to solve.

From the analytical data available at present it seems that the predominant constituent of the pulp is sugar, which, if first converted into alcohol, would place the manufacture of tartaric acid from tamarind on a sound commercial basis, as is done with grapes, first fermenting the juice for the production of alcohol and then utilizing the residue for the manufacture of tartaric acid.

Dr. Sudborough and Mr. P. N. Vridhaachalam say in their paper that although heating in an autoclave is necessary to facilitate the separation of the aqueous extract from the residue, it destroys appreciable amounts of reducing sugars present and the fermentation of the mother liquors from the calcium tartrate precipitate would not be likely to pay.

In order to avoid this loss of sugar, which is a valuable asset in the tamarind pulp, by autoclaving, some preliminary experiments were made in the year 1921 to ferment the pulp extract first, so that the whole of the sugar is made available for the formation of alcohol without undergoing any loss by autoclaving, and then to prepare tartaric acid from the mother liquor after fermentation and also to see the effect of fermentation on the tartaric acid content of the pulp.

Fermented by the organisms, naturally present, the pulp yielded 10.79 per cent. by weight of absolute alcohol, which was determined by ascertaining the specific gravity of the distillates from the fermented must.<sup>1</sup> The same work was simultaneously carried out by Dr. Marsden of Madras, and his findings<sup>2</sup> confirm our results. He reports that 1,000 grm. tamarind were lexiviated, the extract fermented and distilled - giving an yield of alcohol—10.75 per cent. In one of his experiments a yield of 14 per cent. alcohol equivalent to a sugar content of 28 per cent. was also obtained.

Two sets of experiments were started, each in duplicate, to mark the progress of the conversion of sugar into alcohol. The first set was with a pulp containing 33.96 per cent. sugar and the second with a pulp containing 11.60 per cent. sugar. These experiments were carried out in open vats, as is done in distilleries, and the results are given in Tables II and III.

<sup>1</sup> Leffner and Davis. *Allen's Commercial Organic Analysis*, Vol. I.

<sup>2</sup> *Madras Dept. Ind. Bull.* 1 (N.S.).

TABLE II.

*Showing the progress of the conversion of the sugar content of the tamarind pulp into alcohol.*

Days	REDUCING SUGAR PER CENT.		ACTUAL ALCOHOL FORMED PER CENT.		Average actual alcohol formed per cent.	Average theoretical alcohol per cent.	Average loss of alcohol per cent.	Average loss of sugar per cent.
	Vat I	Vat II	Vat I	Vat II				
1	33.96	33.96	..	..	..	..	..	..
2	31.98	31.21	1.05	1.15	1.10	1.21	0.11	0.22
3	25.83	25.83	2.15	3.74	2.94	3.99	0.96	1.88
4	16.18	16.26	7.43	6.03	7.18	9.07	1.89	3.69
5	8.35	9.01	8.03	8.56	8.33	12.87	4.54	8.83
6	5.35	5.73	10.25	9.60	9.93	14.47	4.54	8.83
7	4.85	4.85	10.71	10.79	10.75	14.88	4.13	8.08
8	3.08	3.05	10.17	9.85	10.01	15.79	5.78	11.31

TABLE III.  
*Showing the progress of the conversion of the sugar content of the tamarind pulp into alcohol.*

Days	REDUCING SUGAR PER CENT.		ACTUAL ALCOHOL FORMED PER CENT.		Average theoretical alcohol formed per cent.	Average actual alcohol formed per cent.	Average theoretical alcohol per cent.	Average actual alcohol per cent.	Average loss of sugar per cent.
	Vat I	Vat II	Vat I	Vat II					
1	41.00	41.00	..	..	..	..	..	..	..
2	39.23	39.23	6.78	6.89	6.83	6.83	6.07	6.11	6.11
3	37.44	38.23	1.59	1.59	1.59	1.59	1.62	6.03	6.06
4	37.44	37.44	1.68	1.70	1.69	1.69	1.82	6.13	6.25
5	32.10	32.10	4.10	3.80	3.99	3.99	4.55	6.56	1.60
6	16.65	19.61	11.52	10.48	11.00	11.00	11.84	6.84	1.64
7	6.67	7.13	12.42	11.52	11.97	11.97	17.43	5.46	10.68
8	3.57	3.57	8.47	10.18	9.33	9.33	19.13	9.89	19.17

During fermentation there is a wastage of sugar, which is not a small amount and may be due to the growth of different kinds of bacteria, mucor, etc., in the fermenting liquor. It is also worthy to note here that there is a certain percentage of sugar, though small, which escapes fermentation.

Another experiment was performed in duplicate to see the effect of fermentation on the tartaric acid content of the tamarind pulp. Estimation of tartaric acid was carried out on alternate days and not daily as was the case with sugar in the former experiments. The results are expressed in Table IV.


TABLE IV.

*Showing changes in the tartaric acid content of the tamarind pulp during fermentation.*

Days	TARTARIC ACID PER CENT.		Average tartaric acid per cent.	Average loss of tartaric acid per cent.
	Vat I	Vat II		
1	9.52	9.52	9.52	..
3	9.23	9.20	9.21	0.31
5	9.23	9.23	9.23	0.29
7	8.75	8.99	8.87	0.65
9	8.99	8.87	8.93	0.59

Although there appears to be a little decrease in the quantity of tartaric acid during fermentation, still the outcome of alcohol will considerably outvalue this loss.

The tamarind pulps fermented in the above experiments contained 33.96 and 41.00 per cent. reducing sugars; therefore, the yield of alcohol was even less than  $\frac{1}{3}$ rd the weight of the sugar contents. According to the equation—



the theoretical yield of alcohol is little more than half the quantity of the sugar. In the distilleries of the United Provinces, the parent substance used for the production of alcohol are mostly *mahua* (*Bassia latifolia*) flowers and molasses and the yield of alcohol from each is given in Table V.

TABLE V.

*Showing the yield of alcohol from mahua flowers and molasses.*

Parent substance	Sugar content	Liquor produced	Weight of alcohol per 100 sugar
New mahua flowers . . .	52-55	4½ galls. L.P.	50
Old mahua flowers . . .	42-45	3.8-4 ..	50
Molasses . . . . .	60-62	4½-4 ..	40

The *mahua* flowers give the theoretical yield of alcohol, while molasses gives less, due to the presence of a certain amount of sugar, which escapes fermentation, technically called by distillers as unfermentable sugar.

Thus the amount of alcohol produced from the tamarind is not in proportion to that of the sugar content, as it ought to be. The pulp during fermentation was every now and then examined under a microscope, and it was found that, along with the growth of healthy adventitious yeast cells in strongly acid medium, there was also a growth of different kinds of bacteria, moulds, etc., which may be one of the chief causes of the decreased production of alcohol. Hence it was thought desirable to ferment the pulp with pure cultures of yeasts, which may increase the yield of alcohol. Comparative experiments in duplicate were then carried out on the fermentation of sterilized tamarind must in Erlenmeyer flasks tightly corked with sterilized cotton wool by means of *Saccharomyces Ellipsoides*, *Saccharomyces Exiguus*, *Saccharomyces Capsularis*, beer yeast from Solan Brewery, Simla, yeast from Cawnpore Sugar Works, and adventitious yeasts in tamarind. The results are shown in Table VI.

To make the results easily intelligible the highest figures calculated as percentage of alcohol produced per 100 of tamarind pulp are shown in Plate I. The vertical column represents the percentage of alcohol yielded.

It will thus be seen that *Saccharomyces Ellipsoides* gives a yield of alcohol which nearly approaches the theoretical amount and far exceeds the quantity obtained with other yeasts. The spontaneous fermentation of the must of the tamarind pulp is thus not quite adapted for the production of alcohol.

Besides, during the alcoholic fermentation with adventitious yeast, it was marked in several experiments that the yield of alcohol was



spasmodic and slow, but with pure cultures it was not so. To elucidate this the results of two experiments are shown graphically in Plate II.

TABLE VI.  
*Showing the fermentative efficiency of different species of yeast.*

Name of yeast	Sugar in pulp	ALCOHOL FOUND PER CENT.		Average alcohol found per cent.	Alcohol calcu- lated	Fermen- tative efficiency of yeast per cent.
		Flask I	Flask II			
S. Ellipsoideus . . .	38.87	19.26	19.26	19.26	19.87	96.93
S. Exiguus . . .	38.87	15.93	15.87	15.90	19.87	80.02
S. Capsularis . . .	38.87	13.84	13.76	13.80	19.87	69.45
Beer yeast . . .	38.87	13.86	13.86	13.86	19.87	69.74
Cawnpore yeast . . .	38.87	13.80	13.92	13.86	19.87	69.74
Adventitious yeast . . .	38.87	12.98	12.82	12.90	19.87	64.92


Thus it can be seen that, with *Saccharomyces Ellipsoideus*, there is every day a rise in the yield of alcohol, which was greatest on the 7th day, while with adventitious organisms there was rise and fall, reaching the maximum yield on the 8th day.

In order to find out the influence of the concentration of the must on the production of alcohol, several trials were made in the very beginning to ferment the pulp with varying quantities of water. The fermentation was found to be at its best when the ratio of the pulp to water was between 1 : 3.5 and 1 : 5. Therefore, the above experiments were carried out with one of pulp to five of water at room temperatures which varied from 25° to 39° C. and it was marked that the higher the temperature nearing 40° C. the better and quicker the alcoholic fermentation.


In conclusion, we would, therefore, point out that if the fermentation of the tamarind pulp is to be carried on commercially with pure cultures of yeast in place of adventitious organisms, the yield of alcohol is sure to increase, consequently enhancing the profit calculated by Dr. Marsden for the manufacturers of tartaric acid and alcohol from tamarind.

Finally, we take this opportunity to express our thanks to Dr. and Mrs. Norris of Madras, and Mr. Lissenborough of Cawnpore Sugar Works, for their kindness in supplying us with samples of pure cultures of different varieties of yeasts, and also to B. Manna Lal Khare, Lecturing Assistant, for his help during the investigation.

PLATE I.



# PLATE



## Publications of the Imperial Department of Agriculture in India.

To be had from :—

The Office of the Agricultural Adviser to the Government of India, Pusa, Bihar,  
and from the following Agents :—

- |   |   |
|---|---|
| (1) Thacker, Spink & Co., Calcutta.<br>(2) W. Newman & Co., Calcutta.<br>(3) Rai M. C. Sarkar Bahadur & Sons, Calcutta.<br>(4) Higginbotham & Co., Madras.<br>(5) Thompson, Ltd., Madras. | (6) D. B. Taraporevala, Sons & Co., Bombay.<br>(7) Thacker & Co., Ltd., Bombay.<br>(8) Sunder Pandurang, Bombay.<br>(9) Rai Sahib M. Gulab Singh & Sons, Lahore.<br>(10) Manager, Educational Book Depot, Nagpur. |
|---|---|

**A complete list of the publications of the Imperial Department of Agriculture in India can be obtained on application from the Agricultural Adviser to the Government of India, Pusa, Bihar, or from any of the above-mentioned agents.**

These publications are :—

1. *The Agricultural Journal of India*. A Journal dealing with subjects connected with agricultural economies, field and garden crops, economic plants and fruits, soils, manures, methods of cultivation, irrigation, climatic conditions, insect pests, fungus diseases, co-operative credit, agricultural cattle, farm implements, and other agricultural matters in India. Illustrations, including coloured plates, form a prominent feature of the Journal. It is edited by the Agricultural Adviser to the Government of India, and is issued once every two months or six times a year. *Annual Subscription*, Rs. 6 or 9s. 6d., including postage. Single copy, R. 1-8 or 2s.
2. Scientific Reports of the Agricultural Research Institute, Pusa.
3. Annual Review of Agricultural Operations in India.
4. Proceedings of the Board of Agriculture in India.
5. Proceedings of Sectional Meetings of the Board of Agriculture.
6. Memoirs of the Imperial Department of Agriculture in India :
  - (a) Botanical Series.
  - (b) Chemical Series.
  - (c) Entomological Series.
  - (d) Bacteriological Series.
  - (e) Veterinary Series.
7. Bulletins issued by the Agricultural Research Institute, Pusa.
8. Indigo Publications.
9. Books.

**The following are the publications of the last two years :—**

- Scientific Reports of the Agricultural Research Institute, Pusa (including the Reports of the Imperial Dairy Expert and the Secretary, Sugar Bureau), for the year 1921-22. Price, As. 14.
- Scientific Reports of the Agricultural Research Institute, Pusa (including the Reports of the Imperial Dairy Expert and the Secretary, Sugar Bureau), for the year 1922-23. Price, R. 1.
- Review of Agricultural Operations in India, 1921-22. Price, R. 1-4.
- Review of Agricultural Operations in India, 1922-23. Price R. 1-10.
- Proceedings of the Board of Agriculture in India, held at Bangalore on the 21st January, 1924, and following days (with Appendices). Price, R. 1.
- Proceedings of the Cattle Conference, held at Bangalore on 22nd and 23rd January, 1924 (with appendices). Price, As. 9.
- Report of the Proceedings of the Fifth Entomological Meeting, held at Pusa on 5th—10th February, 1923. Price, Rs. 9-8.
- Proceedings of the Second Meeting of Veterinary Officers in India, held at Calcutta from 26th February to 2nd March, 1923 (with appendices). Price, R. 1-12.

## MEMOIRS OF THE DEPARTMENT OF AGRICULTURE IN INDIA.

### Botanical Series.

- Vol. XI, No. IX. A New Ginger Disease in Godavari District, by S. SUNDARARAMAN, M.A. Price, R. 1 or 1s. 4d.
- Vol. XI, No. X. *Helminthosporium* spp. on Cereals and Sugarcane in India, Part I. Diseases of *Zea Mays* and *Sorghum vulgare* caused by species of *Helminthosporium* by MANORANJAN MITRA, M.Sc. Price, R. 1 or 1s. 4d.
- Vol. XII, No. I. The Wheats of Bihar and Orissa, by ALBERT HOWARD, C.I.E., M.A., GABRIELLE L. C. HOWARD, M.A., and ABDUR RAHMAN KHAN. Price, As. 8 or 7d.
- Vol. XII, No. II. I.—History of the operations against Bud-rot of Palms in South India. II.—Inoculation Experiments with *Phytophthora palmivora* Butl. on *Borassus flabellifer* Linn. and *Cocos nucifera* Linn., by W. McRAE, M.A., B.Sc. Price, R. 1-4 or 2s.
- Vol. XII, No. III. Studies in Inheritance in Cotton, I. History of a Cross between *G. herbaceum* and *G. neglectum*, by G. L. KOTTUR, M.A. Price, R. 1-4 or 1s. 9d.
- Vol. XII, No. IV. Studies in Indian Oil Seeds. No. 2. Linseed, by GABRIELLE L. C. HOWARD, M.A., and ABDUR RAHMAN KHAN. Price, R. 1-4 or 2s.
- Vol. XII, No. V. Studies in Gujarat Cottons, Part II, by MAGANLAL L. PATEL, B.A. Price, R. 1-12 or 3s.
- Vol. XIII, No. I. Studies in Indian Tobaccoes. No. 4.—Parthenocarp and Parthenogenesis in the varieties of *Nicotiana glauca* L. var. *Cuba* and var. *Miradolo*, by GABRIELLE L. C. HOWARD, M.A., and KASHI RAM. No. 5.—The Inheritance of Characters in *Nicotiana rustica* L., by GABRIELLE L. C. HOWARD, M.A. Price, Rs. 2 or 2s. 9d.
- Vol. XIII, No. II. The Wilt Disease of Safflower, by S. D. JOSHI, B.Sc. Price, R. 1 or 1s. 6d.
- Vol. XIII, No. III. Studies in Indian Fibre Plants. No. 3. On the Inheritance of Characters in *Hibiscus Subdariffa* L., by ALBERT HOWARD, C.I.E., and GABRIELLE L. C. HOWARD, M.A. (In the press.)

### Chemical Series.

- Vol. VI, No. VI. Investigations on Indian Opium, No. 3. Studies in the Meconic Acid Content of Indian Opium, by HAROLD E. ARNETT, D.Sc. (Lond.), F.I.C., M.S.E.A.C., and MATHERA NATH BOSE, M.A. Price, As. 6 or 6d.
- Vol. VI, No. VII. Chemical Studies on Safflower Seed and its Germination, by V. A. TAMHANE, M.Sc., M.A. Price, As. 10 or 1s.
- Vol. VI, No. VIII. Note on the Permanent Manurial Plots, Coimbatore, by ROLAND V. NORRIS, D.Sc. Price, R. 1-1 or 1s. 9d.
- Vol. VII, No. I. A Note on the Hydrocyanic Acid in the Burma Bean (*Phaseolus lunatus*, sp.), by F. J. WARTH, M.Sc., B.Sc. Price, As. 12 or 1s.
- Vol. VII, No. II. Studies of an Acid Soil in Assam, II, by A. A. MCGITT, B.Sc. Price, As. 12 or 1s.
- Vol. VII, No. III. A Preliminary Note on the Decomposition of Calcium Cyanamide in South Indian Soils, by ROLAND V. NORRIS, D.Sc., F.I.C., B. VISWANATH and C. V. RAMASWAMI AYYAR, L.A.G. Price, As. 12 or 1s.
- Vol. VII, No. IV. Some Digestibility Trials on Indian Feeding Stuffs, by P. E. LANDER, M.A., D.Sc., A.I.C., and PANDIT LAL CHAND DHARMANI, L.A.G. (In the press.)

### Entomological Series.

- Vol. VII, No. IX. Notes on Indian Diptera, by RONALD SENIOR-WHITE, F.E.S. Price, R. 1-12 or 2s. 3d.
- Vol. VII, No. X. *Platyedra gossypiella*, Saund., Pink Boll-Worm in South India, 1920-21, by E. BALLARD, B.A., F.E.S. Price, As. 12 or 1s.
- Vol. VII, No. XI. Studies in Indian Dermaptera, by MORGAN HERBARD. Price, R. 1-4 or 1s. 9d.
- Vol. VII, No. XII. Further Notes on *Pempheres affinis*, Fst. (The Cotton Stem Weevil), by E. BALLARD, B.A., F.E.S. Price, R. 1 or 1s. 4d.
- Vol. VII, No. XIII. An Account of Experiments on the Control of *Stya (Schani bias) intertellus* in the Godavari Delta, by E. BALLARD, B.A., F.E.S. Price, As. 14 or 1s. 4d.
- Vol. VIII, Nos I, II & III. Hydrophilidæ of India; by A. D'ORCHYMONST; An Annotated List of Ichneumonidæ in the Pusa Collection, by G. R. DUTT, B.A.; A Second Note on Odonata in the Pusa Collection, by MAJOR F. C. FRASER. Price, R. 1 or 1s. 6d.
- Vol. VIII, No. IV. Notes on Indian Muscidæ, by R. SENIOR-WHITE, F.E.S. Price, As. 12 or 1s.

### Veterinary Series.

- Vol. III, No. IV. Studies in Rinderpest, by W. A. POOL, M.B.C.V.S., and T. M. DOYLE, F.R.C.V.S. Price, R. 1 or 1s. 4d.
- Vol. III, No. V. Etiology of Equine Contagious Abortion in India, by T. M. DOYLE, F.R.C.V.S. Price, As. 8 or 9d.

### BULLETINS ISSUED BY THE AGRICULTURAL RESEARCH INSTITUTE, PUSA.

- No. 136. The Hydrogen Ion Concentrations of some Indian Soils and Plant Juices, by W. R. C. ATKINS, O.B.E., Sc.D., F.R.C. Price, As. 4.
- No. 137. Note on the Probability of an Inter-relation between the Length of the Stigma and that of the Fibre in some forms of the genus *Gossypium*, by RAM PRASAD. Price, As. 4.
- No. 138. Methods of Examination of Certain Characters in Cotton, by G. R. HULSON, B.Sc. Price, As. 8.
- No. 139. List of Publications of Indian Entomology, 1920-21 (compiled by the Imperial Entomologist). Price, R. 1.
- No. 140. The Determination of Prussic Acid in Burma Beans (*Phaseolus lunatus*) (Preliminary Note), by J. CHARLTON, B.Sc. Price, As. 3.
- No. 141. Comparative Manurial Value of the whole Plants and the different parts of Green Manures, by N. V. JOSHI, B.A., M.Sc., L.A.G. Price, As. 6.
- No. 142. The Cultivation of Lac in the Plains of India, by C. S. MISRA, B.A. Price, R. 1-8.
- No. 143. *Ber (Zizyphus jujuba)* Fruit and its Fly Pest, by J. L. KHARE, F.E.S. Price, As. 6.
- No. 144. Observations on the Morphology and Life-cycle of *Filaria recandita* Grassi, by M. ANANT NARAYAN RAO. Price, As. 6.
- No. 145. Some Observations on the Barren Soils of Lower Bari Doab Colony in the Punjab, by S.M. NASIR. Price, As. 3.
- No. 146. The Prevention of Nuisances caused by the Parboiling of Paddy, by J. CHARLTON, B.Sc., A.L.C. Price, As. 5.
- No. 147. List of Publications on Indian Entomology, 1922 (compiled by the Imperial Entomologist). Price, As. 7.
- No. 148. The Relative Responsibility of Physical Heat and Micro-organisms for the Hot Weather Rotting of Potatoes in Western India, by S. L. AJERKAR, B.A., and J. D. RANADIVE, B.A.G. Price, As. 5.

**BULLETINS ISSUED BY THE AGRICULTURAL RESEARCH INSTITUTE,  
PUSA—conold.**

- No. 149. A Study of the Factors operative in the value of Green Manure, by B. H. WILSDON, B.A., L.E.S., P. E. LANDER, M.A., D.Sc., A.L.C., and M. MUKAND LAL. Price, As. 5.
- No. 150. The Improvement of Fodder and Forage in India (Papers read before a joint meeting of the sections of Agriculture and Botany, Indian Science Congress, 1923), edited by GABRIELLE L. C. HOWARD, M.A. Price, As. 6.
- No. 151. A Method for the Accurate Determination of Carbonic Acid present as Carbonate in Soils, by PHANI BHUSAN SANYAL, M.Sc. Price, As. 2.
- No. 152. The External Morphology and Bionomics of the Commonest Indian Tick (*Hyalomma aegyptium*), by MOHAMMAD SHARIF, M.A., F.R.M.S. Price, R. 1.
- No. 153. Tamarind as a source of Alcohol and Tartaric Acid, by H. N. Batham, M.A., and L. S. Nigam, L.A.G. Price, As. 3 or 4d.
- No. 154. The Bionomics of the Sarcopic Mange Parasite of the Buffalo with some observations concerning the relative power of resistance to adverse conditions of the different stages of the Acarus and of its egg, by T. M. TIMONEY, M.R.C.V.S. Price, As. 2.
- No. 155. List of Publications on Indian Entomology, 1923 (compiled by the Imperial Entomologist). (*In the press.*)

---

**INDIGO PUBLICATIONS.**

- No. 11. The Nature of Changes Occurring in the Indigo Steeping Vat, II, by W. A. DAVIS, B.Sc., A.C.S.I. Price, R. 1-10.
- No. 12. Indigo Experiments, 1922. (1) The effect on produce when vat liquor is allowed to stand in the beating vat and beating is delayed. (2) The effect of neutralizing the liquor with caustic soda before beating, by J. H. WALTON, M.A., M.Sc. Price, As. 4.

---

**BOOKS.**

- Wheat in India, by ALBERT HOWARD, M.A., A.R.C.S., F.I.S., and GABRIELLE L. C. HOWARD, M.A. Price, Rs. 5 or 7s. 6d.
- A Description of the Imperial Bacteriological Laboratory, Muktesar: Its Work and Products, by Major J. D. E. HOLMES, M.A., D.Sc., M.R.C.V.S. Price, As. 8 or 9d.
- Agriculture in India, by JAMES MACKENNA, M.A., L.C.S. Price, As. 4 or 5d.
- Some Diseases of Cattle in India. A Handbook for Stock-owners. Price, As. 8 or 9d.
- The Importance of Bacterial Action in Indigo Manufacture, by C. M. HUTCHINSON, B.A. Price, As. 2 or 3d.
- Report on the Diseases of Silkworms in India, by A. PRINGLE JAMESON, D.Sc. Price, Rs. 3.

---

**NOTICE.**

All new publications are included in the above list and are regularly posted to all on the mailing list as soon as issued. Unless complaints of non receipt of any publication are received within six months of the date of issue, duplicate copies can be supplied only on payment of cost and postage.

